

Effects of Optigen® in commercial dairy herd diets on milk composition, milk production, and economics

JF INOSTROZA¹, RD SHAVER¹, VE CABRERA¹, AND JM TRICÁRICO²

¹Department of Dairy Science, University of Wisconsin, Madison, WI, USA ²Alltech Inc., Brookings, SD, USA

Introduction

While nutritionists are typically concerned with the effect of protein source on ration cost, there is a growing appreciation of the cost of inefficient crude protein (CP) usage. As a result, nonprotein nitrogen (NPN) is used in dairy rations as a less expensive alternative to protein of plant or animal origin.

Objectives

To determine the effect of Optigen® (blended, controlled-release urea), as a source of dietary nitrogen on milk composition, milk production, and economics in commercial dairy herds in Wisconsin.

Materials and Methods

Animals and experimental design

- 16 commercial Wisconsin dairy herds (average 148 cows/herd; range 58 – 550 cows/herd).
- Crossover experimental design using two 30-d feeding periods.
- Within herds, cows were fed a single-diet TMR.
- Across the 16 herds, TMR contained 56±3% forages composed of 43±9% corn silage and formulated for 17.1±0.4% CP and 30.5±1.7% NDF (DM basis).

Treatments

- **Control** – TMR
- **Optigen®** – TMR with 114 g/cow/d Optigen® replacing an equivalent amount of supplemental CP, primarily from SBM.
- Treatments were isonitrogenous.
- Diet formulation space created by Optigen® was filled with DM from either corn grain or corn silage.

Measurements

- Weight and composition (fat, protein, MUN) of bulk tank milk shipments per herd were recorded.
- Number of cows with milk in the bulk tank per shipment was recorded.
- Average daily milk yield and milk component yields were calculated.
- A spreadsheet economic simulation was performed using the Optigen® feeding rate and milk yield response from the field trial and monthly (as-fed) prices for SBM-48 (\$0.373±0.054/kg), dry corn (\$0.188 ±0.020/kg), corn silage (\$0.059±0.005/kg), and high-moisture corn (\$0.149 ±0.016/kg) and milk prices (\$0.38±0.03/kg) for January – December, 2008. Local pricing for Optigen® was used. A total of 32 combinations of varying feed and milk prices were simulated.

Data analysis

- Mixed model procedure of SAS with period, sequence and treatment as fixed effects and herd as a random effect.

Results

- Milk yield was 0.5 kg/d/cow greater (P<0.01) for Optigen® compared with the control (Table 1).
- Optigen® reduced feed cost only when corn silage was used to fill the formulation space. However, milk income minus feed cost was increased by Optigen® for all forms of dry corn used to fill the formulation space. This difference was greatest for the corn silage treatment at \$0.21/cow/d.
- A decision tool spreadsheet was developed to allow additional economic simulations. The tool allowed for varying the following: milk yield response to Optigen®, cost of Optigen®, and the CP and energy supplements evaluated (Table 2).

Table 1. Milk production and milk composition in response to Optigen®.

	Control	Optigen®	SEM	P-Value
Milk yield, kg/d	35.4	35.9	0.2	0.01
Fat, %	3.72	3.69	0.02	0.07
yield, g/d	1317	1322	8	NS
Protein, %	2.98	2.97	0.01	NS
yield, g/d	1055	1065	6	0.13
MUN, mg/dL	12.4	13.2	0.3	0.01

Table 2. Economic impact of Optigen® use in dairy herd diets.

Crude protein supplement replaced by Optigen®	Ingredient used to fill formulation space	Feed cost OPT - CON (\$/cow/d)	Milk income OPT - CON (\$/cow/d)	Milk income minus feed cost (\$/cow/d)
SBM-48	Dry corn	0.047 (± 0.027)	0.192 (± 0.016)	0.145 (± 0.039)
SBM-48	Corn silage	-0.020 (± 0.039)	0.192 (± 0.016)	0.212 (± 0.051)
SBM-48	High-moisture corn	0.042 (± 0.028)	0.192 (± 0.016)	0.150 (± 0.040)

Conclusions

- Under the conditions of this study, Optigen® reduced feed cost only when corn silage was used to fill the formulation space, however milk income minus feed cost was increased by Optigen® for all forms of corn used.
- A decision tool spreadsheet was developed to help producers, consultants, and extension agents compare and select feed supplements in diets of lactating cows.

Acknowledgements

The author thanks his advisor Randy Shaver for the support with the trial design, statistical analysis and very useful discussions. Also thanks to Victor Cabrera for his contribution to develop the Decision Tool. Appreciation is expressed to Todd LeNoble and Jennifer Meyer, from CP Feeds, for their contribution with information for some dairies that participated in the study; to Spence Driver, John Weinkes, Randy Greenfield, Rob Sjoerdsma, Mike Yahr and Irv Haugen, from Vitaplus Company, for their contribution with information for dairies that we used in the trial. Finally, thanks to all the Wisconsin dairy farmers that contributed with this study.